



AMERICAN
SPEECH-LANGUAGE-
HEARING
ASSOCIATION

Updated Evidence-Based Systematic Review: Effects of Intensity of Treatment and Constraint-Induced Language Therapy for Individuals With Stroke-Induced Aphasia

Leora Cherney
Rehabilitation Institute of Chicago, Chicago, IL

Janet Patterson
California State University, Hayward, CA

Anastasia Raymer
Old Dominion University, Norfolk, VA

Tobi Frymark and Tracy Schooling
National Center for Evidence-Based Practice in Communication Disorders,
American Speech-Language-Hearing Association, Rockville, MD

Background

This is an updated version of the original evidence-based systematic review (EBSR) published in 2008 in the *Journal of Speech, Language, and Hearing Research* (see <http://jslhr.asha.org/cgi/content/full/51/5/1282>).

The use of constraint-induced language treatment (CILT) by speech-language pathologists continues to gain momentum as a viable treatment for individuals with stroke-induced aphasia. Since our original systematic review of the literature (Cherney, Patterson, Raymer, Frymark, & Schooling, 2008), additional research has emerged investigating the key principles of CILT—*forced use of verbal language* and *high-intensity massed practice* (Pulvermüller et al., 2001).

Objective

To evaluate the effects of intensity and CILT on individuals with stroke-induced aphasia. See Appendix A1 for the original clinical questions.

Search strategy

An updated search of 24 electronic databases such as MEDLINE (January 2006 to August 2010), PsycINFO (January 2006 to August 2010), *The Cochrane Library* (February issue, 2010), and CINAHL (January 2006 to August 2010) was conducted using key words related to stroke-induced aphasia, amount and intensity of treatment, and CILT. The titles and abstracts of all articles were independently assessed by two reviewers and the full texts of relevant articles were obtained for further scrutiny. References of all full-text articles were scanned to identify additional literature of relevance. See Appendix A2 for search methodology.

Selection criteria

Inclusion required meeting the following criteria: (a) English-only studies published in peer-reviewed journals with original data addressing one or more of the original clinical questions, (b) active treatment examining CILT or directly comparing two different intensities of an aphasia treatment, and (c) included participants ages 18 years of age or older with stroke-induced acute or chronic aphasia. Exclusion criteria consisted of studies pertaining to individuals with cognitive deficits as well as mixed treatments or populations in which data could not be analyzed separately. Studies that used pharmacological intervention for aphasia treatment were also excluded. See Appendix A3 for the list of excluded studies and reasons for exclusion.

Data collection and analysis

Two reviewers independently applied the inclusion criteria, assessed study quality and extracted data from included studies using ASHA's levels of evidence scheme outlined in the previous version of this review (Cherney et al., 2008).

Main results

In the original 2008 review, we identified 10 studies relevant to the treatment of stroke-induced aphasia, with five studies investigating treatment intensity (see <http://jslhr.asha.org/cgi/content/full/51/5/1282/T7>), four studies examining CILT (see <http://jslhr.asha.org/cgi/content/full/51/5/1282/T8>), and one examining both. This updated review identified 16 new studies—13 pertaining to CILT and three related to intensity of aphasia treatment.

Table 1 displays the clinical questions addressed, stage of research, and methodological quality of the included studies. Of these, only three (Bakheit et al., 2007; Goral & Kempler, 2009; Meinzer, Streiftau, & Rockstroh, 2007) were considered efficacy research. The remaining 13 were classified as exploratory research. Similar to the original review, the majority of studies described participants and treatments sufficiently to allow for study replication. All but two (Breier, Maher, Schmadeke, Hasan, & Papanicolaou, 2007; Harnish, Neils-Strunjas, Lamy, & Eliassen, 2008) provided probability data and 13 of 16 (81%) allowed for calculation of treatment effects for one or more reported outcomes.

Table 2 provides a detailed description of the 202 participants studied, including type, severity, and chronicity of aphasia. Of the participants included, half (50.5%; 102/202) exhibited chronic aphasia with time post onset (TPO) ranging from 6 months to 12 years and half (49.5%; 100/202) exhibited acute aphasia with TPO \leq 3 months (Bakheit et al., 2007, treatment intensity; Kirmess & Maher, 2010, CILT). Thirteen studies reported aphasia classification type (fluent vs. nonfluent) or type of aphasia (e.g., Broca's, Wernicke's, conduction). The majority of studies included participants with nonfluent aphasia ranging from moderate to severe.

CILT results: Table 3 outlines the treatment variables and major findings of the 13 studies examining CILT. One hundred of the 202 total participants received CILT, with 4% (4/100) at the acute phase of recovery and 96% (96/100) at the chronic phase of recovery. All 13 studies contributed data to examine the effects of CILT on measures of language impairment. Six also examined the effects of CILT on measures of communication activity/participation (Breier, Maher, Novak, & Papanicolaou, 2006; Brier et al., 2007, 2009; Faroqi-Shah & Virion, 2009; Goral & Kempler, 2009; Kirmess & Maher, 2010) and four (Breier et al., 2007; Faroqi-Shah & Virion, 2009; Goral & Kempler, 2009; Kirmess & Maher, 2010) addressed the maintenance effects of treatment after CILT. Interestingly, the exploratory studies revealed mixed results in support of CILT, while the two efficacy studies (Goral & Kempler, 2009; Meinzer, Streiftau, & Rockstroh, 2007) reported statistically significant changes favoring CILT on various measures of language impairment and communication activity/participation. It is important to note that several included CILT studies have modified the technique from the original description of Pulvermüller et al. (2001), but the key principles of constraint, massed practice, and shaping were retained.

Intensity results: Table 4 illustrates the treatment variables and major findings of the 102 participants from three studies investigating the effect of treatment intensity on measures of language impairment, the majority of which (95%; 97 of 102) came from the Bakheit et al. (2007) trial. This study compared standard care (2 hr per week) to intensive treatment (5 hr per week) for individuals receiving speech and language intervention at the acute stage of recovery. No significant difference was found between more intensive ($M = 35.6$ hr) and less intensive ($M = 19.3$ hr) treatment on the Western Aphasia Battery Aphasia Quotient (Kertesz, 1982). The remaining two studies addressed language impairment outcomes at the chronic stage of recovery. Of these studies, only one (Ramsberger & Marie, 2007) provided statistical data to compare a nonintensive (2 times per week) to an intensive (5 times per week) computer naming treatment. The authors reported mixed findings, with two of the four participants favoring more intensive treatment and two favoring less intensive treatment. Although all four participants demonstrated maintenance of treatment gains, only one (Participant 3) was considered significant. These

findings differ from the original review in which the majority of studies found increased treatment intensity to be associated with positive changes in outcome measures of language impairment.

Authors' conclusions

While the update of this review uncovered additional literature supporting the benefits of CILT for individuals with stroke-induced aphasia, the conclusion and clinical message remains unchanged. The majority of the evidence supporting the use of CILT is exploratory in nature. Future research beyond the exploratory phase of study is warranted to determine the efficacy of CILT. Given the findings from Bakheit et al. (2007), further review of studies that provide high intensity massed practice, an important component of CILT, should be considered.

Finally, although not specific to our clinical questions, it is important to note that a number of studies included in this review provided neurophysiologic data examining the neurocorrelates of behavioral changes. Investigation focusing on changes in neural activation patterns will further contribute to our understanding of CILT.

At this time, although research findings suggest the positive effects of CILT, clinicians should interpret the findings of this review judiciously, and as always, in conjunction with clinical expertise and the patient's preferences to determine the most appropriate treatment plan.

Acknowledgements

This work was supported by the American Speech-Language-Hearing Association's National Center for Evidence-Based Practice in Communication Disorders.

References

- Bakheit, A. M., Shaw, S., Barrett, L., Wood, J., Carrington, S., Griffiths, S., ... Koutsi, F. (2007). A prospective, randomized, parallel group, controlled study of the effect of intensity of speech and language therapy on early recovery from poststroke aphasia. *Clinical Rehabilitation, 21*, 885–894.
- Breier, J. I., Juranek, J., Maher, L. M., Schmadeke, S., Men, D., & Papanicolaou, A. C. (2009). Behavioral and neurophysiologic response to therapy for chronic aphasia. *Archives of Physical Medicine & Rehabilitation, 90*, 2026–2033.
- Breier, J. I., Maher, L. M., Novak, B., & Papanicolaou, A. C. (2006). Functional imaging before and after constraint-induced language therapy for aphasia using magnetoencephalography. *Neurocase, 12*, 322–331.
- Breier, J. I., Maher, L. M., Schmadeke, S., Hasan, K. M., & Papanicolaou, A. C. (2007). Changes in language-specific brain activation after therapy for aphasia using magnetoencephalography: A case study. *Neurocase, 13*, 169–177.
- Cherney, L., Patterson, J., Raymer, S., Frymark, T., & Schooling, T. (2008). Evidence-based systematic review: Effects of intensity of treatment and constraint-induced language therapy for individuals with stroke-induced aphasia. *Journal of Speech, Language, and Hearing Research, 51*, 1282–1299.
- Faroqi-Shah, Y., & Virion, C. R. (2009). Constraint-induced language therapy for agrammatism: Role of grammaticality constraints. *Aphasiology, 23*, 977–988.
- Goral, M., & Kempler, D. (2009). Training verb production in communicative context: Evidence from a person with chronic non-fluent aphasia. *Aphasiology, 23*, 1383–1397.
- Harnish, S. M., Neils-Strunjas, J., Lamy, M., & Eliassen, J. C. (2008). Use of fMRI in the study of chronic aphasia recovery after therapy: A case study. *Topics in Stroke Rehabilitation, 15*, 468–483.
- Kertesz, A. (1982). *Western Aphasia Battery*. Orlando, FL: Grune & Stratton.
- Kirmess, M., & Maher, L. (2010). Constraint induced language therapy in early aphasia rehabilitation. *Aphasiology, 24*, 725–736.
- Meinzer, M., Fleisch, T., Breitenstein, C., Wienbruch, C., Elbert, T., & Rockstroh, B. (2008). Functional re-recruitment of dysfunctional brain areas predicts language recovery in chronic aphasia. *Neuroimage, 39*, 2038–2046.
- Meinzer, M., Fleisch, T., Obleser, J., Assadollahi, R., Djundja, D., Barthel, G., & Rockstroh, B. (2006). Brain regions essential for improved lexical access in an aged aphasic patient: A case report. *BioMedCentral Neurology, 6*, 28.
- Meinzer, M., Obleser, J., Fleisch, T., Eulitz, C., & Rockstroh, B. (2007). Recovery from aphasia as a function of language therapy in an early bilingual patient demonstrated by fMRI. *Neuropsychologia, 45*, 1247–1256.
- Meinzer, M., Paul, I., Wienbruch, C., Djundja, D., & Rockstroh, B. (2009). Electromagnetic brain activity in higher frequency bands during automatic word processing indicates

- recovery of function in aphasia. *European Journal of Physical Rehabilitation & Medicine*, 45, 369–378.
- Meinzer, M., Streiftau, S., & Rockstroh, B. (2007). Intensive language training in the rehabilitation of chronic aphasia: Efficient training by laypersons. *Journal of International Neuropsychological Society*, 13, 846–853.
- Pulvermüller, F., Neininger, B., Elbert, T., Mohr, B., Rockstroh, B., Koebbel, P., & Taub, E. (2001). Constraint-induced therapy of chronic aphasia after stroke. *Stroke*, 32, 1621–1626.
- Ramsberger, G., & Marie, B. (2007). Self-administered cued naming therapy: A single-participant investigation of a computer-based therapy program replicated in four cases. *American Journal of Speech-Language Pathology*, 16, 343–358.
- Richter, M., Miltner, W. H., & Straube, T. (2008). Association between therapy outcome and right-hemispheric activation in chronic aphasia. *Brain*, 131, 1391–1401.
- Szaflarski, J. P., Ball, A., Grether, S., Al-fwaress, F., Griffith, N. M., Neils-Strunjas, J., ... Reichhardt, R. (2008). Constraint-induced aphasia therapy stimulates language recovery in patients with chronic aphasia after ischemic stroke. *Medical Science Monitor*, 14, 243–250.

Table 1. Methodological quality of included studies, stage of research, and clinical question(s) addressed.

| Citation | Question(s) research phase | Study design | Protocol description | Group/subject comparability | Assessor blinding | Sampling allocation | Treatment fidelity | Significance | Precision | ITT | Quality indicators |
|---------------------------------------|----------------------------|----------------|----------------------|-----------------------------|-------------------|--------------------------------|--------------------|--------------|-----------|-----|--------------------|
| | | | | | | | | | | | |
| Bakheit et al., 2007 (Intensity) | Q 3 Efficacy | Trial | + | + | + | Random; adequate description | - | + | + | + | |
| Breier et al., 2006 (CILT) | Q 6,7 Exploratory | Case series | + | + | - | Convenience | + | + | + | N/A | |
| Breier et al., 2007 (CILT) | Q 6,7,10 Exploratory | Case study | + | + | - | Convenience | - | - | - | N/A | |
| Breier et al., 2009 (CILT) | Q 6,7 Exploratory | Case series | + | + | - | Convenience | - | + | + | N/A | |
| Faroqi-Shah & Virion, 2009 (CILT) | Q 6,7,10 Exploratory | Case series | + | +/- | + | Convenience | - | + | + | N/A | |
| Goral & Kempfer, 2009 (CILT) | Q 6,7,10 Efficacy | Single subject | + | + | + | Convenience | - | + | + | N/A | |
| Harmish et al., 2008 (Intensity) | Q 1 Exploratory | Case study | + | + | - | Convenience | - | - | - | N/A | |
| Kirmess & Maher, 2010 (CILT) | Q 8,9,10 Exploratory | Case series | + | + | - | Convenience | + | + | + | N/A | |
| Meinzer et al., 2006 (CILT) | Q 6 Exploratory | Case study | + | + | - | Convenience | - | + | - | N/A | |
| Meinzer et al., 2007 (CILT) | Q 6 Exploratory | Case study | + | + | - | Convenience | - | + | - | N/A | |
| Meinzer et al., 2008 (CILT) | Q 6 Exploratory | Case series | + | + | - | Convenience | - | + | + | N/A | |
| Meinzer et al., 2009 (CILT) | Q 6 Exploratory | Case series | + | + | - | Convenience | - | + | + | N/A | |
| Meinzer, Streiftau, & Rockstroh, 2007 | Q 6 Efficacy | Trial | + | + | + | Random; inadequate description | - | + | + | + | |

| (CILT) | | | | | | | | | | | | |
|--|----------------------|-------------------|---|---|---|---|-----------------------------------|---|---|---|---|-----|
| Ramsberger & Marie, 2007 (Intensity) | Q 1,5 Exploratory | Single subject | + | + | - | + | Random; inadequate description | + | + | + | + | N/A |
| Richter, Miltner, & Siraube, 2008 (CILT) | Q 6 Exploratory | Case series | + | + | - | - | Convenience | - | + | + | + | N/A |
| Szaflarski et al., 2008 (CILT) | Q 6,7 Exploratory | Case series | + | + | - | - | Convenience | - | + | + | + | N/A |

Note. ITT = intention to treat analysis; N/A = not applicable; Q = question.

Table 2. Participant characteristics.

| Study | N | Age (years) | Education (years) | Gender M:F | RH | TPO (years) | Etiology | Aphasia type | Aphasia severity | Aphasia test score |
|-----------------------------------|----|---|--------------------------|----------------|------------|--------------------------------------|---------------|---------------------------|------------------|---|
| Bakheit et al., 2007 (Intensity) | 97 | Intensive: 71.2 (14.9) Standard: 69.7 (15.0) | 10.9 (2.9) 10.5 (2.1) | 26:25 21:25 | 90% 93% | 34.2 days (19.1) 28.1 days (14.9) | 85 TE 12 H | — | — | I WAB AQ M = 44.2 (30.2) SC WAB AQ M = 37.9 (27.2) |
| Breier et al., 2006 (CILT) | 6 | 62.0 (9.7) | 13.5 (1.9) | 4:2 | 100% | 3.9 (1.7) | 6 I | 5 Broca's 1 conduction | Mod-severe | WAB AQ M = 52.2 (22.0) |
| Breier et al., 2007 (CILT) | 1 | 62.0 | 16 | F | 100% | 1.1 | H | — | Mod-severe | WAB AQ = 43.0 |
| Breier et al., 2009 (CILT) | 23 | 54.0 (11.0) | 12+ (0.08) | 16:7 | 100% | ≥1 | 2 H 21 I | — | Mod-severe | WAB AQ M = 57.0 (17.0) BNT M = 19.0 (17.0) |
| Faroqi-Shah & Virion, 2009 (CILT) | 4 | 57.8 (9.8) | 16.8 (3.0) | 3:1 | 100% | 5.1 (4.5) | — | 3 Broca's 1 agrammatic | — | WAB AQ M = 75.5 (13.2) BNT M = 45.8 (12.4) OAN M = 38.8 (5.1) |
| Goral & Kempler, 2009 (CILT) | 1 | 60.0 | — | M | — | 12 | — | Nonfluent | — | CLQT CS = 3.4 BDAE-3 Aud. Comp. = 12%ile BDAE-3 Cookie Theft = 13%ile BDAE-3 Repetition = 55%ile BDAE-3 Naming = 37%ile BDAE-3 Reading = 26%ile BDAE-3 Writing = 18%ile |
| Harmish et al., 2008 (Intensity) | 1 | 52.0 | — | F | — | 8 | — | Conduction | — | WAB AQ = 76.1 |
| Kirness & | 3 | 66.7 (23.0) | >12 | 2:1 | 100% | 46.8 days (9.86) | 1 H | Nonfluent | 2 severe 1 NR | NGA overall = 110.3 |

2 NR

Maher,
2010
(CILT)

| | | | | | | | | | | |
|---|----|---|------------|------|------|------------------------|-------------|--|-----------------------------|---|
| Meinzer et al., 2006 (CILT) | 1 | 80.0 | — | F | 100% | 2.1 | I | Wernicke's | — | AAT PS = 47.5 |
| Meinzer et al., 2007 (CILT) | 1 | 35.0 | 18+ | M | 100% | 2.7 | I | Amnestic | Mod | — |
| Meinzer et al., 2008 (CILT) | 11 | 49.1 (13.4) | — | 7;4 | 100% | 6.0 (11.3) | 2 H 9 I | 7 Broca's 1 global 2 Wernicke's 1 not classified | 4 mild 6 mod 1 severe | AAT PS M = 51.8 (4.5) |
| Meinzer et al., 2009 (CILT) | 10 | 45.1 (17.5) | 10.7 (1.8) | 4;6 | 100% | 0.6 (1.5) | 2 H 8 I | 1 amnestic 4 Broca's 2 global 1 Wernicke's 1 not classified | 3 mild 7 mod | AAT PS M = 50.1 (5.5) |
| Meinzer, Streifflau, & Rockstroh, 2007 (CILT) | 20 | CILT-SLP: 50.2 (10.1) CILT-R: 62 (8.9) | 11 (1.6) | 16;4 | 100% | 2.6 (1.6) 3.9 (1.4) | 2 H 18 I | 2 amnestic 11 Broca's 3 global 3 Wernicke's 1 not classified | 8 mild 9 mod 3 severe | SLP AAT PS M = 50.6 (4.8) R AAT PS M = 52.7 (4.0) |
| Ramsberger & Marie, 2007 (Intensity) | 4 | 67.5 (5.4) | 12+ | 1;3 | — | 2.6 (2.4) | — | 1 anomic 1 Broca's 1 Wernicke's 1 conduction | — | PI WAB AQ = 69 P2 WAB AQ = 53 P3 ADP SS = 90 P4 ADP SS = 107 |
| Richter, Milner, & Straube, 2008 (CILT) | 16 | 58.3 (9.6) | — | 12;4 | 100% | ≥1 | — | 7 anomic 7 Broca's 2 global | — | AAT SS M = 3.3 (0.91) AAT TT M = 33.8 (16.7) |
| Szaflarski et al., 2008 (CILT) | 3 | 58-64 | 14.6 | 3;0 | 100% | ≥2 | 3 I | 1 fluent 2 nonfluent | Mod-severe | BDAE-3 Aud. Comp. = 28.9 BDAE-3 Verbal Exp M = 25 |

Note. Dashes indicate data not reported by study author. ADP SS = Aphasia Diagnostic Profile standard score; AAT PS = Aachen Aphasia Test; Aud. Comp. = auditory comprehension; BDAE = Boston Diagnostic Aphasia Examination; BNT = Boston Naming Test; CILT = constraint induced language treatment; CILT-R = constraint induced language treatment–relative; CILT-SLP = constraint induced language treatment–speech-language pathologist; CLQT CS= Cognitive Linguistic Quick Test-Composite Score; H = hemorrhagic; I = ischemic; NGA = Norwegian Basic Aphasia Assessment; PS = Profile Score of the Aachen Aphasia Test; SS = Speech Scale of the Aachen Aphasia Test; TPO = time post onset, TT = Token Test of the Aachen Aphasia Test; Verbal Exp = verbal expression; WAB AQ = Western Aphasia Battery aphasia quotient.

Table 3. CILT treatment variables and findings

| Study | Treatment program/schedule | Outcome type | Outcome measure(s) | Effect size | p |
|---------------------|---|---------------------|---------------------------|---------------------|----------|
| Breier et al., 2006 | <i>Treatment:</i> CILT principles Dual card task with visual barrier Dyad of participants <i>Schedule:</i> 3 hr per day 4 days per week for 3 weeks <i>Total:</i> 36 hr | I | WAB | | |
| | | | AQ | 0.9 [-1.04, 1.22] | .228 |
| | | | Repetition | 0.00 [-1.13, 1.13] | 1.0 |
| | | | Auditory Comprehension | 0.03 [-1.10, 1.17] | .59 |
| | | | BNT | -0.03 [-1.16, 1.10] | .738 |
| | | | % CIUs | | |
| | | | Dual card probes | 0.57 [-0.59, 2.25] | .022* |
| Accuracy task | 1.04 [-0.16, 1.72] | .088 | | | |
| Breier et al., 2007 | <i>Treatment:</i> CILT principles Dual card task with visual barrier Dyad of participant & clinician <i>Schedule:</i> 3 hr per day 4 days per week for 3 weeks <i>Total:</i> 36 hr | I | WAB | | |
| | | | AQ | | |
| | | | Information Content | | |
| | | | Fluency | | |
| | | | Auditory Comprehension | | |
| | | | Repetition | | |
| | | | Naming | | |
| | | | BNT | | |
| | | | PALPA | | |
| | | | Word-Picture Matching | | |
| | | | Sentence-Picture Matching | | |
| A/P | % CIUs | | | | |

| | | | | | | |
|----------------------------|--|-----|---|---|--------------------------------|--------|
| | | | Dual card task probes | — | — | — |
| | | | Generalization untrained probes | — | — | — |
| Breier et al., 2009 | <i>Treatment: CILT principles</i> Dual card task with visual barrier Dyad of participants <i>Schedule: 3 hr per day</i> 4 days per week for 3 weeks <i>Total: 36 hr</i> | I | WAB | | | |
| | | A/P | AQ | — | — | <.004* |
| | | | % CIUs | | | |
| | | | Dual card task (<i>N</i> = 9) | 1.06 [0.08, 2.05] | | <.05* |
| Faroqi-Shah & Virion, 2009 | <i>Treatment: CILT principles</i> Dual card task with visual barrier & additional tense morphology shaping Dyad not stated <i>Schedule: 3 hr per day for 10 days</i> <i>Total: 24 hr</i> | I | WAB + BNT + OAB (language <i>t</i> score) | Post: 0.18 [-1.21, 1.57] <i>f/u: 0.38 [-1.01, 1.78]</i> | Post: .075 <i>f/u: .127</i> | |
| | | I | WAB AQ | Post: -0.76 [-0.67, 2.2] <i>f/u: -0.09 [-1.48, 1.29]</i> | Post: .178 <i>f/u: .498</i> | |
| | | I | OAB | Post: 0.29 [-1.10, 1.68] <i>f/u: 0.84 [-0.61, 2.28]</i> | Post: .65 <i>f/u: .353</i> | |
| | | I | BNT | Post: -0.21 [1.61, 1.18] <i>f/u: 0.38 [-1.02, 1.77]</i> | Post: .348 <i>f/u: .074</i> | |
| | | I | Verb Inflection Test | Post: 0.84 [-0.61, 2.28] <i>f/u: -0.37 [-1.77, 1.03]</i> | Post: .066 <i>f/u: .08</i> | |
| | | A/P | Cinderella Narrative & Conv. | | | |
| | | | Proportion Sentences | Post: 0.00 [-1.39, 1.39] <i>f/u: -0.27, -1.66, 1.12]</i> | Post: .9 <i>f/u: .493</i> | |
| | | | Proportion of Well Formed Sentences | Post: 0.61 [-0.8, 2.03] <i>f/u: -0.11 [-1.49, 1.28]</i> | .059 <i>f/u: .884</i> | |
| | | | Tense Accuracy | Post: 0.4 [-1.0, 1.8] <i>f/u: -0.05 [-1.44, 1.34]</i> | Post: .364 <i>f/u: .962</i> | |
| | | | Tense Diversity | Post: 0.75 [-0.44, 2.25] <i>f/u: 0.44 [-0.96, 1.85]</i> | Post: .192 <i>f/u: .444</i> | |
| Goral & Kempler, 2009 | <i>Treatment: CILT principles</i> Dual card task with visual barrier | I | BDAE | — | — | — |
| | | A/P | Treatment Block 1 | | | |

| | | | | | |
|--------------------------|--|-----------|--|--------------------------------------|-----------|
| Kirmess & Maher, 2010 | Structured tasks with target verbs (e.g., picture description). Unstructured tasks with no target verb (e.g., story construction). 2-3 tasks per session. Dyad of participant & clinician <i>Schedule:</i> 5 hr per day for 4 days <i>Total:</i> 20 hr per treatment block | A/P | Narrative – Total Words | Post: < 1 f/u: -1.96 | <i>ns</i> |
| | | | Narrative – % of Verbs | Post: 1.97 ^a f/u: < -1 | <i>ns</i> |
| | | | Treatment Block 2 | | |
| | | | Narrative – Total Words | Post: 8 f/u: < 1 | — |
| | | | Narrative – % of Verbs | Post: 2.2 f/u: < -1 | — |
| | | | Social Communication | 0.43 [-0.21, 1.05] | .003* |
| | | | NGA | | |
| | | | Overall | 0.34 [-1.23, 2.00] | — |
| | | | Naming | 0.85 [-0.09, 2.42] | .030* |
| | | | Auditory Comprehension | 0.03 [-1.58, 1.62] | .789 |
| Meinzer et al., 2006 | <i>Treatment:</i> CILT principles Dual card task with visual barrier Dyad of participants <i>Schedule:</i> 1.15-3 hr per day for 10 days <i>Total:</i> M = 24.8 hr | I | Writing | 0.32 [-1.35, 1.89] | .547 |
| | | | PALPA Naming Frequency | 0.75 [-1.18, 2.06] | .167 |
| | | | VOST Sentence Construction | 1.16 [-0.95, 2.34] | .138 |
| | | | TROG-2 | 0.12 [-1.47, 1.74] | .141 |
| | | | CILT | | |
| Meinzer et al., 2006 | <i>Treatment:</i> CILT principles Dual card task with visual barrier Dyad not stated <i>Schedule:</i> 3 hr per day for 10 days <i>Total:</i> 30 hrs | I | Dual card task – high & low frequency requests | 0.93 [-1.10, 2.16] | — |
| | | | AAT | | |
| | | | Profile Score | — | S |
| | | | Token Test – Errors | — | <i>ns</i> |
| | | | Repeating | — | S |
| Written Language | — | <i>ns</i> | | | |

| | | | | |
|---|-------------------------|--------------------------|--------------------|-----------|
| | | Naming | — | <i>ns</i> |
| | | Comprehension | — | <i>ns</i> |
| I | | Naming Test | | |
| | | Corrected Responses | — | — |
| | | Self-Corrected Responses | — | — |
| I | Meinzer et al., 2007 | AAT | | |
| | | Profile Score | — | S |
| | | Token Test - Errors | — | S |
| | | Naming | — | <i>ns</i> |
| | | Naming Test | — | — |
| I | | AAT | | |
| | | Profile Score | 0.45 [-0.40, 1.30] | <.0001* |
| | | Token Test – Errors | 0.31 [-0.53, 1.15] | .003* |
| | | Repeating | 0.21 [-0.63, 1.04] | .002* |
| | | Written Language | 0.31 [-0.33, 1.15] | .009* |
| | | Naming | 0.34 [-0.50, 1.19] | .004* |
| | | Comprehension | 0.22 [-0.62, 1.06] | .008* |
| I | | Naming | | |
| | | Trained Items | 1.11 [0.21, 2.01] | .0008* |
| | | Untrained Items | 0.62 [-0.23, 1.48] | .002* |
| I | Meinzer et al., 2009 | AAT | | |
| | | Profile Score | 0.34 [-0.55, 1.22] | <.001* |
| | | Written Language | 0.23 [-0.65, 1.11] | <.02* |
| I | Meinzer, Streiftau, | AAT – SLP | | |

& Rockstroh,
2007

Dual card task with visual barrier
Dyad of participants
Schedule: 3 hours per day for 10 days
Total: 30 hr

| | | |
|------------------|--------------------|---------|
| Profile Score | 0.45 [-0.44, 1.33] | <.0001* |
| Token Test | 0.31 [-0.57, 1.19] | <.008* |
| Repeating | 0.51 [-0.38, 1.40] | <.004* |
| Written Language | 0.18 [-0.70, 1.06] | <.036* |
| Naming | 0.31 [-0.57, 1.20] | <.01* |
| Comprehension | 0.31 [-0.57, 1.19] | <.009* |
| AAT – Relatives | | |

I

| | | |
|--------------------|--------------------|---------|
| Profile Score | 0.54 [-0.35, 1.44] | <.0002* |
| Token Test | 0.39 [-0.49, 1.28] | <.0002* |
| Repeating | 0.53 [-0.36, 1.43] | <.0006* |
| Written Language | 0.20 [-0.68, 1.08] | <.006* |
| Naming | 0.46 [-0.43, 1.34] | <.014* |
| Comprehension | 0.48 [-0.41, 1.36] | <.002* |
| Relatives vs. SLPs | | |

I

| | | |
|------------------|---------------------|------|
| Profile Score | -0.45 [-1.34, 0.44] | >.27 |
| Token Test | 0.15 [-0.73, 1.03] | >.77 |
| Repeating | -0.84 [-1.75, 0.08] | >.46 |
| Written Language | 0.00 [-1.41, 1.41] | >.17 |
| Naming | -0.36 [-1.25, 0.52] | >.59 |
| Comprehension | 0.43 [-0.46, 1.32] | >.77 |

Richter, Miltner,
& Straube, 2008

Treatment: CILT principles
Dual card task with visual barrier
Dyad of participants
Schedule: 3 hr per day for 10 days
Total: 30 hr

| | | |
|----------------------------|---------------------|-----------|
| AAT | | |
| Token Test | -0.05 [-0.76, 0.65] | <i>ns</i> |
| Spontaneous Speech | 0.37 [-0.34, 1.08] | <.0001* |
| ANELT | | |
| Auditory Comprehensibility | 0.28 [-0.42, 0.91] | <.01* |

I

| | | Semantic Comprehensibility | 0.59 [-0.12, 1.29] | <.05* | |
|----------------------------|--|----------------------------|---------------------------------|---------------------|------|
| Szaflarski et al., 2008 | <i>Treatment:</i> CILT principles Dual card task with visual barrier Dyad of participants <i>Schedule:</i> 3 hr per day plus 30–45 min informal socialization breaks for 5 days <i>Total:</i> 15 hr | I | BDAE-3 | | |
| | | | Auditory Comprehension | 0.27 [-1.34, 1.87] | .328 |
| | | | Verbal Expression | -0.04 [-1.64, 1.56] | .423 |
| | | | Language Communication Index | 0.14 [-1.47, 1.74] | .312 |
| | | | Fable Retell – Utterances | 0.80 [-1.24, 2.83] | .090 |
| | | | Fable Retell – Words | 0.31 [-1.66, 2.28] | .051 |
| | | | Fable Retell – Word Roots | 0.99 [-1.09, 3.07] | .186 |
| | | | Fable Retell – MLU | 0.02 [-1.94, 1.98] | .947 |
| | | | Fable Retell – TTR | 0.16 [-1.8, 2.13] | .618 |

Note. Dashes indicate data not reported by study author or data not calculable. ANELT = Amsterdam Nijmegen Everyday Language Test; AAT = Aachen Aphasia Test; A/P = activity/participation measure; AQ = aphasia quotient; BDAE-3 = Boston Diagnostic Aphasia Examination-3; BNT = Boston Naming Test; CIUs = content information units; I = impairment measure; MLU = mean length of utterance; NGA = Norwegian Basic Aphasia Assessment; OAB = Object Action Battery; PALPA = Psycholinguistic Assessment of Language Processing in Aphasia; S = significant; TROG-2 = Test for Reception of Grammar-2; TTR = type-token ratio; VOST = Verb and Sentence Test; WAB = Western Aphasia Battery.

^aEffect size change based on Goral & Kempler reported correction.

*Denotes a significant change.

Table 4. Intervention variables and findings for intensity studies.

| Study | Treatment program/schedule | Outcome type | Outcome measure(s) | Effect size | <i>p</i> |
|--------------------------|---|--------------|--------------------|-------------------------|---------------------------|
| Bakheit et al., 2007 | <i>Treatment:</i> targeting naming and conversation <i>Intensive:</i> 5 hr per week; <i>M</i> = 35.6 hr (16.4); total sessions: 37.16 (14.9) <i>Standard Care:</i> 2 hr per week; <i>M</i> = 19.3 hr (6.4); total sessions: 8.6 (6.4) <i>Total:</i> 12 weeks | I | WAB AQ | | |
| | | | Week 4 | 0.05 [-0.37, 0.48] | .88 |
| | | | Week 8 | 0.09 [-0.37, 0.54] | .98 |
| | | | Week 12 | 0.15 [-0.31, 0.61] | <i>ns</i> |
| | | | Week 24 | 0.07 [-0.41, 0.56] | <i>ns</i> |
| Harnish et al., 2008 | <i>Treatment:</i> targeting word finding and phonological processing <i>Massed:</i> 1.5 hr per day; 5 days per week for 2 weeks <i>Distributive:</i> 1 hr per day; 2 days per week for 7.5 weeks <i>Total:</i> 15 hr massed; 15 hr distributive | I | BNT | | |
| | | | Massed | — | — |
| | | | Distributive | — | — |
| Ramsberger & Marie, 2007 | <i>Treatment:</i> MossTalk computer naming treatment 45–60 min sessions <i>Nonintensive:</i> 2 sessions per week; total: 15 sessions <i>Intensive:</i> 5 sessions per week; total: 20 sessions | I | Naming | | |
| | | | Participant 1 | | |
| | | | Intensive | Post: 5.53 f/u: 3.61 | Post: .03* f/u: .26 |
| | | | Nonintensive | Post: 4.00 | Post: <.01* |
| | | | Participant 2 | | |
| | | | Intensive | Post: 2.08 f/u: 2.20 | Post: .07 f/u: .06 |
| | | | Nonintensive | Post: 2.44 | Post: .09 |
| | | | Participant 3 | | |
| | | | Intensive | Post: 4.68 f/u: 4.70 | Post: <.01* f/u: <.01* |
| | | | Nonintensive | Post: 4.98 | Post: <.01* |
| Participant 4 | | | | | |
| Intensive | Post: 3.79 | Post: <.01* | | | |

| | | |
|--------------|------------|------------|
| | f/u: 5.74 | f/u: .06 |
| Nonintensive | Post: 4.96 | Post: .03* |

Note. Dashes indicate not reported by study author or not calculable. A/P = activity/participation; BNT = Boston Naming Test; I = impairment; WAB AQ= Western Aphasia Battery aphasia quotient.

*Denotes a significant change.

Appendix

A1. Clinical questions

Intensity:

1. For stroke-induced chronic aphasia, what is the influence of treatment intensity on measures of language impairment?
2. For stroke-induced chronic aphasia, what is the influence of treatment intensity on measures of communication activity/participation?
3. For stroke-induced acute aphasia, what is the influence of treatment intensity on measures of language impairment?
4. For stroke-induced acute aphasia, what is the influence of treatment intensity on measures of communication activity/participation?
5. For stroke-induced chronic aphasia, what treatment outcomes are maintained following intensive language treatment?

CILT:

6. For stroke-induced chronic aphasia, what is the influence of constraint-induced language therapy on measures of language impairment?
7. For stroke-induced chronic aphasia, what is the influence of constraint-induced language therapy on measures of communication activity/participation?
8. For stroke-induced acute aphasia, what is the influence of constraint-induced language therapy on measures of language impairment?
9. For stroke-induced acute aphasia, what is the influence of constraint-induced language therapy on measures of communication activity/participation?
10. For stroke-induced chronic aphasia, what treatment outcomes are maintained following constraint-induced language therapy?

A2. Search Methodology

Electronic Databases Searched:

PubMed (www.ncbi.nlm.nih.gov/pubmed/)
CINAHL (EBSCO)
Health Source: Nursing (EBSCO)
Psychology and Behavioral Sciences Collection (EBSCO)
PsycINFO (EBSCO)
Communication & Mass Media Complete (EBSCO)
Education Research Complete (EBSCO)
ComDisDome (CSA)
CSA Neurosciences Abstracts
ERIC (CSA)
LLBA (CSA)
CSA Social Services Abstracts
ISI Web of Science
Cochrane Library (Wiley)
ScienceDirect
SUMSearch (<http://sumsearch.uthscsa.edu/>)
SpeechBITE (www.speechbite.com/)
REHABDATA (www.naric.com/research/rehab/)
TRIP Database (www.tripdatabase.com/)
Latin American and Caribbean Center on Health Sciences Information (LILACS)
(www.bireme.br/php/index.php?lang=en)
NHS Evidence Health Information Resources (formerly National Library for Health)
(www.library.nhs.uk/)
The Aphasiology Archive (<http://aphasiology.pitt.edu/>)
HighWire Press (<http://highwire.stanford.edu>)
GoogleScholar

Search Criteria:

- Date range: 2006 to present
- Peer-reviewed journal (including in-press studies)
- English only
- Must be a study with original data that addresses one or more of the clinical questions
- For clinical questions related to intensity, the keywords *intensity* or *amount of treatment* must be included in the abstract or title and studies must directly compare frequency/intensity of treatment.

Inclusion Criteria:

- Adults ages 18 and older
- Individuals with stroke-induced aphasia

Exclusion Criteria:

- Studies with individuals with cognitive deficits
- Studies that include pharmacological intervention
- Individual studies that are included in accepted systematic reviews or meta-analyses

Search Terms:

Constraint Induced Language Treatment OR CILT

Constraint Language Treatment OR CLT

Constraint Induced Aphasia Treatment OR CIAT

Constraint Aphasia Treatment

Forced Language Treatment

Intensive Language-Action Therapy OR ILAT

Aphasia (exploded MeSH term)

Stroke (exploded MeSH term)

Amount OR Intensity OR Frequency OR Duration OR Length of Treatment

Treatment Schedule

Verbal Expression or Speech Output

Adult

Expanded Search Terms:

("Aphasia/rehabilitation"[Mesh] OR "Aphasia/therapy"[Mesh]) AND ("Stroke"[Mesh]) AND ("Treatment Outcome"[Mesh] OR amount OR intense OR intensit* OR frequent OR frequenc* OR duration OR length OR schedule OR session)

("Aphasia/rehabilitation"[Mesh] OR "Aphasia/therapy"[Mesh]) AND ("Language Therapy/methods*"[Mesh])

(constrain* induced aphasia)

(constrain* OR forced OR intensive OR intense OR intensity) AND (language OR aphasia) AND (treatment OR therapy OR program)

"Cerebrovascular disorders"[MeSH Major Topic] AND "aphasia/therapy"[MeSH Major Topic]

"Cerebrovascular disorders"[MeSH Major Topic] AND "aphasia "[MeSH Major Topic] AND (amount OR intens* OR freq* OR duration OR length OR schedule)

"Aphasia/therapy"[MeSH Major Topic] AND stroke

"Aphasia/rehabilitation"[MeSH Major Topic] AND (stroke OR "cerebrovascular disorders"[MeSH Major Topic])

"Aphasia"[MeSH Major Topic] AND stroke AND (amount OR intens* OR freq* OR duration OR length OR schedule)

"Aphasia/rehabilitation"[MeSH Major Topic] OR "aphasia/therapy"[MeSH Major Topic]

(Constrain* OR forced) AND “language therapy”

CILT OR CIAT OR CLT

“Language”[MeSH Major Topic] AND (constrain* OR forced) AND (treatment OR therapy OR rehabilitation OR intens*)

“Aphasia/therapy”[MeSH Major Topic] AND “time factors”[MeSH Major Topic]
("Cerebrovascular disorders"[MeSH Terms] OR "cerebrovascular accident"[MeSH Terms])
AND "rehabilitation of speech and language disorders"[MeSH Terms]

("Cerebrovascular disorders/rehabilitation"[MeSH Terms] OR "cerebrovascular
accident/rehabilitation"[MeSH Terms]) AND (speech OR language OR communication OR
aphasi*)

("Cerebrovascular disorders/rehabilitation"[MeSH Terms] OR "cerebrovascular
accident/rehabilitation"[MeSH Terms]) AND ("language disorders"[MeSH Terms] OR "speech
disorders/rehabilitation"[MeSH Terms])

"Aphasia"[MeSH Terms] AND (amount OR intens* OR freq* OR duration OR length OR
schedule)

Model orientated aphasia therapy

MH = Major Heading

RH = Rehabilitation

TH = Therapy

(MH "Aphasia/RH/TH") AND (constrain* OR forced OR intensive OR intense OR intensity)

(MH "Aphasia/RH/TH") AND (amount OR intensit* OR frequent OR frequenc* OR duration
OR length OR schedule OR session)

(MH "Stroke/RH/TH") AND (aphasi* OR language OR constrain* OR forced OR verbal OR
speech) AND (amount OR intensit* OR frequent OR frequenc* OR duration OR length OR
schedule OR session)

(Constrain* induced language) OR (constrain* language) OR forced language OR forced
language treatment

((MH "Cerebrovascular Disorders+")) AND (MH "Aphasia+/RH/TH")

((MH "Cerebral Vascular Accident") OR (MH "Cerebral Vascular Accident")) AND (MH
"Rehabilitation, Speech and Language+")

(MH "Aphasia+/RH/TH") AND (amount OR intens* OR freq* OR duration OR treatment schedule)

((MH "Cerebrovascular Disorders+") OR (MH "Cerebral Vascular Accident")) AND (MH "Language Disorders+") and (MH "Rehabilitation, Speech and Language+")

(MH "Aphasia/RH/TH") AND stroke

(MH "Aphasia+") AND (amount OR intens* OR freq* OR duration OR length OR schedule)

((MH "Cerebrovascular Disorders+") OR (MH "Cerebral Vascular Accident")) AND aphasia

DE= Subject

(DE "APHASIA" AND PT "ACADEMIC JOURNAL") OR (DE "APHASIA --*" AND PT "ACADEMIC JOURNAL")

(constrain* induced aphasia)

(constrain* OR forced OR intensive OR intense OR intensity) AND (language OR aphasia) AND (treatment OR therapy OR program)

(model orientated aphasia therapy) OR (intensive language action therapy)

(constrain* OR forced OR intensive OR intense OR intensity) AND aphasia treatment

DE = Thesaurus term

(DE "Aphasia" OR DE "Acalculia" OR DE "Agnosia" OR DE "Agraphia" OR DE "Dysphasia") AND (language OR constrain* OR forced OR verbal OR speech) AND (amount OR intensit* OR frequent OR frequenc* OR duration OR length OR schedule OR session)

(DE "APHASIA" OR DE "AGRAMMATISM" OR DE "ANOMIA" OR DE "CONDUCTION aphasia" OR DE "JARGON aphasia" OR DE "WORD deafness") AND (language OR constrain* OR forced OR verbal OR speech) AND (amount OR intensit* OR frequent OR frequenc* OR duration OR length OR schedule OR session)

((Aphasia AND stroke) OR (aphasia AND (amount OR intens* OR freq* OR duration OR length OR schedule)) OR (constrain* AND (language OR aphasia))) AND DE "APHASIA"

“treatment of aphasia” exploded

DE=descriptor

DE=("aphasia 03400") AND DE=("language therapy 44400")
DE="language therapy" and DE="stroke rehabilitation"

Q1=article subject terms

Q1="language" and Q1="aphasia" AND (constrain* OR forced OR amount OR intensive OR intensit* OR frequent OR frequenc* OR duration OR length OR schedule OR session)

DE=descriptor

DE="aphasia" and DE="outcomes of treatment" and DE="therapy"

(DE "Aphasia") AND stroke

(DE "Aphasia") AND (amount OR intens* OR freq* OR duration OR length OR schedule)

(Stroke) AND (DE "Time Factors (Learning)")

(DE "Aphasia") AND (DE "Time Factors (Learning)")

(Stroke) AND (amount OR intens* OR freq* OR duration OR length OR schedule)

(Stroke) AND (DE "Communication Disorders")

(DE "Communication Disorders") AND ((DE "Time Factors (Learning)") OR (amount OR intens* OR freq* OR duration OR length OR schedule))

(Constrain* induced language/aphasia) OR (constrain * language/aphasia) OR forced language/aphasia

DE="aphasia 03400" and DE="language therapy 44400"

DE=("aphasia" AND "language therapy") AND (stroke OR (amount OR intens* OR freq* OR duration OR length OR schedule))

DE="language therapy" AND (amount OR intens* OR freq* OR duration OR length OR schedule) AND stroke

DE=("aphasia" AND "language therapy") AND (speak* OR speech OR spoken OR verbal OR express*)

DE=("aphasia") AND (treatment OR therapy OR rehabilitation) and (amount OR intens* OR freq* OR duration OR length OR schedule)

(DE="language therapy" OR "speech therapy") AND (speak* OR speech OR spoken OR verbal OR express*) AND (amount OR intens* OR freq* OR duration OR length OR schedule)

(constrain* induced aphasia) OR (intensive language action therapy)

(constrain* OR forced OR intensive OR intense OR intensity) AND (language OR aphasia)
AND (treatment OR therapy OR program)

(constrain* OR forced OR intensive OR intense OR intensity) AND (aphasia treatment)

(constrain* OR forced) AND (language therapy)
("Aphasia/rehabilitation"[Mesh] OR "Aphasia/therapy"[Mesh]) AND ("Stroke"[Mesh]) AND
("Treatment Outcome"[Mesh] OR amount OR intense OR intensit* OR frequent OR frequenc*
OR duration OR length OR schedule OR session)

("Aphasia/rehabilitation"[Mesh] OR "Aphasia/therapy"[Mesh]) AND ("Language
Therapy/methods*"[Mesh])

Anywhere in Text: constraint induced language

Anywhere in Text: constraint induced aphasia

Anywhere in Text: constraint constrained forced intensive intense intensity
Title & Abstract only: language aphasia

Title & Abstract only: intensive language action therapy

Anywhere in Text: constraint constrained forced intensive intense intensity
Title & Abstract only: language therapy

Target area: aphasia
Intervention: language therapy
Client sub-groups: stroke/CVA
Age group: adults

Intensive language
Age group: adults

Keyword(s): forced :: Target area: Aphasia

Keyword(s): intensity :: Target area: Aphasia

aphasia [Abstract words] and constraint [Abstract words]

aphasia [Abstract words] and constrained [Abstract words]

aphasia [Abstract words] and forced [Abstract words]

aphasia [Abstract words] and intensity [Abstract words] and treatment [Abstract words]
aphasia [Abstract words] and intense [Abstract words] and treatment [Abstract words]

aphasia [Abstract words] and intensive [Abstract words] and treatment [Abstract words]
language [Abstract words] and intensity [Abstract words] and treatment [Abstract words]
language [Abstract words] and intense [Abstract words] and treatment [Abstract words]
language [Abstract words] and intensive [Abstract words] and treatment [Abstract words]
ciat [Abstract words] or cilt [Abstract words] or clt [Abstract words]

"constraint induced aphasia"

"constraint induced language"

aphasia AND intens* AND (treatment OR therapy)

Additional Searches:

- The reference lists of all relevant articles identified were scanned for other possible studies.
- All accepted articles were forward-searched on ISI Web of Science.
- All ASHA journals were searched through the HighWire Press website.
- Contemporary Issues in Communication Science and Disorders (CICSD) was hand searched.

The literature search was conducted from March to June 2010 by Beverly Wang. References were managed using the bibliographic software EndNote.

A3. Bibliography of excluded studies

Reason for exclusion: could not separate data from mixed treatments or mixed populations

1. Breier, J. I., Randle, S., Maher, L. M., & Papanicolaou, A. C. (2010). Changes in maps of language activity activation following melodic intonation therapy using magnetoencephalography: Two case studies. *Journal of Clinical and Experimental Neuropsychology*, 32(3), 309–314. Advance online publication. doi: 10.1080/13803390903029293
2. Kavanagh, D. O., Lynam, C., Duerk, T., Casey, M., & Eustace, P. W. (2010). Variations in the presentation of aphasia in patients with closed head injuries. *Case Reports in Medicine*, 5.
3. Martin, P. I., Naeser, M. A., Ho, M., Treglia, E., Kaplan, E., Baker, E. H., & Pascual-Leone, A. (2009). Research with transcranial magnetic stimulation in the treatment of aphasia. *Current Neurology and Neuroscience Reports*, 9, 451–458.
4. Seniów, J., Litwin, M., Litwin, T., Leśniak, M., & Członkowska, A. (2009). New approach to the rehabilitation of post-stroke focal cognitive syndrome: Effect of levodopa combined with speech and language therapy on functional recovery from aphasia. *Journal of the Neurological Sciences*, 283(1–2), 214–218.

Reason for exclusion: pharmacological treatment

5. Berthier, M. L., Green, C., Lara, J. P., Higuera, C., Barbancho, M. A., Dávila, G., & Pulvermüller, F. (2009). Memantine and constraint-induced aphasia therapy in chronic poststroke aphasia. *Annals of Neurology*, 65, 577–585.

Reason for exclusion: did not address clinical question

6. Abel, S., Willmes, K., & Huber, W. (2007). Model-oriented naming therapy: Testing predictions of a connectionist model. *Aphasiology*, 21, 411–447.
7. Ansaldo, A. I., Marcotte, K., Scherer, L., & Raboyeau, G. (2008). Language therapy and bilingual aphasia: Clinical implications of psycholinguistic and neuroimaging research. *Journal of Neurolinguistics*, 21, 539–557.
8. Archibald, L. M. D., Orange, J. B., & Jamieson, D. J. (2009). Implementation of computer-based language therapy in aphasia. *Therapeutic Advances in Neurological Disorders*, 2(5), 299–311.
9. Bakheit, A., Shaw, S., Carrington, S., & Griffiths, S. (2007). The rate and extent of improvement with therapy from the different types of aphasia in the first year after stroke. *Clinical Rehabilitation*, 21, 941–949.
10. Bayley, M. T., Hurdowar, A., Teasell, R., Wood-Dauphinee, S., Korner-Bitensky, N., Richards, C. L., ... Jutai, J. W. (2007). Priorities for stroke rehabilitation and research: Results of a 2003 Canadian Stroke Network consensus conference. *Archives of Physical Medicine and Rehabilitation*, 88, 526–528.

11. Beeke, S., Maxim, J., & Wilkinson, R. (2007). Using conversation analysis to assess and treat people with aphasia. *Seminars in Speech and Language, 28*(2), 136–147.
12. Brunner, M., Skeat, J., & Morris, M. E. (2008). Outcomes of speech-language pathology following stroke: Investigation of inpatient rehabilitation and rehabilitation in the home programs. *International Journal of Speech-Language Pathology, 10*, 305–313.
13. Burns, M. S. (2008). Application of neuroscience to technology in stroke rehabilitation. *Topics in Stroke Rehabilitation, 15*, 570–579.
14. Carter, A. R., Connor, L. T., & Dromerick, A. W. (2010). Rehabilitation after stroke: Current state of the science. *Current Neurology and Neuroscience Reports, 10*(3), 158–166.
15. Cherney, L., Small, S., Wier, H., Lobo, N., & Erickson, R. (2006). Functional magnetic resonance imaging changes in brain activity after intensive language therapy with and without cortical stimulation in chronic Broca's aphasia. *Archives of Physical Medicine and Rehabilitation, 87*(10), E5.
16. Cherney, L. R. (2010). Oral reading for language in aphasia: Impact of aphasia severity on cross-modal outcomes in chronic nonfluent aphasia. *Seminars in Speech and Language, 31*(1), 42–51.
17. Cherney, L. R., Halper, A. S., Holland, A. L., & Cole, R. (2008). Computerized script training for aphasia: Preliminary results. *American Journal of Speech-Language Pathology, 17*(1), 19–34.
18. Cherney, L. R., Halper, A. S., Holland, A. L., Lee, J. B., Babbitt, E., & Cole, R. (2007). Improving conversational script production in aphasia with virtual therapist computer treatment software. *Brain & Language, 103*(1–2), 246–247.
19. Cherney, L. R., & Small, S. L. (2006). Task-dependent changes in brain activation following therapy for nonfluent aphasia: Discussion of two individual cases. *Journal of the International Neuropsychological Society, 12*, 828–842.
20. Code, C., Torney, A., Gildea-Howardine, E., & Willmes, K. (2010). Outcome of a one-month therapy intensive for chronic aphasia: Variable individual responses. *Seminars in Speech and Language, 31*(1), 21–33.
21. Conroy, P., Sage, K., & Ralph, M. A. L. (2006). Towards theory-driven therapies for aphasic verb impairments: A review of current theory and practice. *Aphasiology, 20*, 1159–1185.
22. Corsten, S., Mende, M., Cholewa, J., & Huber, W. (2007). Treatment of input and output phonology in aphasia: A single case study... 36th Clinical Aphasiology Conference, Ghent University, Ghent, Belgium, May 29th to June 2nd, 2006. *Aphasiology, 21*(6–8), 587–603.
23. Crosson, B., McGregor, K., Gopinath, K. S., Conway, T. W., Benjamin, M., Chang, Y. L., ... White, K. D. (2007). Functional MRI of language in aphasia: A review of the literature and the methodological challenges. *Neuropsychology Review, 17*(2), 157–177.
24. Davis, C. H., Harrington, G., & Baynes, K. (2006). Intensive semantic intervention in fluent aphasia: A pilot study with fMRI. *Aphasiology, 20*(1), 59–83.

25. del Toro, C. M., Altmann, L. J. P., Raymer, A. M., Leon, S., Blonder, L. X., & Gonzalez Rothi, L. J. (2008). Changes in aphasic discourse after contrasting treatments for anomia. *Aphasiology*, *22*(7–8), 881–892.
26. Edmonds, L. A., Nadeau, S. E., & Kiran, S. (2009). Effect of verb network strengthening treatment (VNeST) on lexical retrieval of content words in sentences in persons with aphasia. *Aphasiology*, *23*(3), 402–424.
27. Faroqi-Shah, Y. (2008). A comparison of two theoretically driven treatments for verb inflection deficits in aphasia. *Neuropsychologia*, *46*, 3088–3100.
28. Fox, S., Armstrong, E., & Boles, L. (2009). Conversational treatment in mild aphasia: A case study. *Aphasiology*, *23*(7–8), 951–964.
29. Fridriksson, J., Morrow-Odom, L., Moser, D., Fridriksson, A., & Baylis, G. (2006). Neural recruitment associated with anomia treatment in aphasia. *Neuroimage*, *32*(3), 1403–1412.
30. Fridriksson, J., Moser, D., Bonilha, L., Morrow-Odom, K. L., Shaw, H., Fridriksson, A., ... Rorden, C. (2007). Neural correlates of phonological and semantic-based anomia treatment in aphasia. *Neuropsychologia*, *45*, 1812–1822.
31. Fridriksson, J., Nettles, C., Davis, M., Morrow, L., & Montgomery, A. (2006). Functional communication and executive function in aphasia. *Clinical Linguistics & Phonetics*, *20*, 401–410.
32. Grande, M., Hussmann, K., Bay, E., Christoph, S., Piefke, M., Willmes, K., & Huber, W. (2008). Basic parameters of spontaneous speech as a sensitive method for measuring change during the course of aphasia. *International Journal of Language and Communication Disorders*, *43*, 408–426.
33. Greener, J., Enderby, P., & Whurr, R. (2001). Pharmacological treatment for aphasia following stroke. *Cochrane Database of Systematic Reviews* (4), CD000424.
34. Hu, M. H., Hsu, S. S., Yip, P. K., Jeng, J. S., & Wang, Y. H. (2010). Early and intensive rehabilitation predicts good functional outcomes in patients admitted to the stroke intensive care unit. *Disability and Rehabilitation*, *32*, 1251–1259.
35. Jordan, L. C., & Hillis, A. E. (2006). Disorders of speech and language: aphasia, apraxia and dysarthria. *Current Opinion in Neurology*, *19*, 580–585.
36. Karges, J., & Smallfield, S. (2009). A description of the outcomes, frequency, duration, and intensity of occupational, physical, and speech therapy in inpatient stroke rehabilitation. *Journal of Allied Health*, *38*(1), E1–10.
37. Kelly, H., & Armstrong, L. (2009). New word learning in people with aphasia. *Aphasiology*, *23*, 1398–1417.
38. Kendall, D. L., Nadeau, S. E., Conway, T., Fuller, R. H., Riestra, A., & Gonzalez Rothi, L. J. (2006). Treatability of different components of aphasia—insights from a case study. *Journal of Rehabilitation, Research and Development*, *43*, 323–336.
39. Kendall, D. L., Rodriguez, A. D., Rosenbek, J. C., Conway, T., & Gonzalez Rothi, L. J. (2006). Influence of intensive phonomotor rehabilitation on apraxia of speech. *Journal of Rehabilitation, Research and Development*, *43*, 409–418.

40. Kiran, S., Sandberg, C., & Abbott, K. (2009). Treatment for lexical retrieval using abstract and concrete words in persons with aphasia: Effect of complexity. *Aphasiology*, *23*(7–8), 835–853.
41. Knollman-Poter, K. (2008). Acquired apraxia of speech: A review. *Topics in Stroke Rehabilitation*, *15*, 484–493.
42. Koenig-Bruhin, M., & Studer-Eichenberger, F. (2007). Therapy of short-term memory disorders in fluent aphasia: A single case study. *Aphasiology*, *21*, 448–458.
43. Laganaro, M., Di Pietro, M., & Schnider, A. (2006). Computerised treatment of anomia in acute aphasia: Treatment intensity and training size. *Neuropsychological Rehabilitation*, *16*, 630–640.
44. Laska, A. C., Kahan, T., Hellblom, A., Murray, V., & von Arbin, M. (2008). Design and methods of a randomized controlled trial on early speech and language therapy in patients with acute stroke and aphasia. *Topics in Stroke Rehabilitation*, *15*, 256–261.
45. Lee, J. B., Kaye, R. C., & Cherney, L. R. (2009). Conversational script performance in adults with non-fluent aphasia: Treatment intensity and aphasia severity. *Aphasiology*, *23*, 885–897.
46. Lee, J., Fowler, R., Rodney, D., Cherney, L., & Small, S. L. (2010). IMITATE: An intensive computer-based treatment for aphasia based on action observation and imitation. *Aphasiology*, *24*, 449–465.
47. Manheim, L. M., Halper, A. S., & Cherney, L. (2009). Patient-reported changes in communication after computer-based script training for aphasia. *Archives of Physical Medicine and Rehabilitation*, *90*, 623–627.
48. Marini, A., Caltagirone, C., Pasqualetti, P., & Carlomagno, S. (2007). Patterns of language improvement in adults with non-chronic non-fluent aphasia after specific therapies. *Aphasiology*, *21*, 164–186.
49. Menke, R., Meinzer, M., Kugel, H., Deppe, M., Baumgartner, A., Schiffbauer, H., ... Breitenstein, C. (2009). Imaging short- and long-term training success in chronic aphasia. *BMC Neuroscience*, *10*, 118.
50. Moss, A., & Nicholas, M. (2006). Language rehabilitation in chronic aphasia and time postonset: A review of single-subject data. *Stroke*, *37*, 3043–3051.
51. Price, C. J., Seghier, M. L., & Leff, A. P. (2010). Predicting language outcome and recovery after stroke: The PLORAS system. *Nature Reviews Neurology*, *6*, 202–210.
52. Raboyeau, G., De Boissezon, X., Marie, N., Balduyck, S., Puel, M., Bézy, C., ... Cardebat, D. (2008). Right hemisphere activation in recovery from aphasia: Lesion effect or function recruitment? *Neurology*, *70*, 290–298.
53. Raymer, A., & Kohen, F. (2006). Word-retrieval treatment in aphasia: Effects of sentence context. *Journal of Rehabilitation, Research and Development*, *43*, 367–378.
54. Raymer, A., Strobel, J., Prokup, T., Thomason, B., & Reff, K. L. (2010). Errorless versus errorful training of spelling in individuals with acquired dysgraphia. *Neuropsychological Rehabilitation*, *20*(1), 1–15.

55. Raymer, A. M., Ciampitti, M., Holliway, B., Singletary, F., Blonder, L. X., Ketterson, T., ... Gonzalez Rothi, L. J. (2007). Semantic-phonologic treatment for noun and verb retrieval impairments in aphasia. *Neuropsychological Rehabilitation, 17*(2), 244–270.
56. Raymer, A. M., Singletary, F., Rodriguez, A., Ciampitti, M., Heilman, K. M., & Rothi, L. J. (2006). Effects of gesture+verbal treatment for noun and verb retrieval in aphasia. *Journal of the International Neuropsychological Society, 12*, 867–882.
57. Rijntjes, M. (2006). Mechanisms of recovery in stroke patients with hemiparesis or aphasia: New insights, old questions and the meaning of therapies. *Current Opinion in Neurology, 19*(1), 76–83.
58. Schlaug, G., Marchina, S., & Norton, A. (2009). Evidence for plasticity in white-matter tracts of patients with chronic Broca's aphasia undergoing intense intonation-based speech therapy. *Annals of the New York Academy of Sciences, 1169*, 385–394.
59. Schröder, C., Schupp, W., Seewald, B., & Haase, I. (2007). Computer-aided therapy in aphasia therapy: Evaluation of assignment criteria. *International Journal of Rehabilitation Research, 30*, 289–295.
60. Small, S. L., Cherney, L. R., Erickson, R. K., Wier, H. Y., & Lobo, N. C. (2006). Intensive language therapy with and without cortical stimulation in nonfluent aphasia. *Annals of Neurology, 60*(Suppl. 3), S61.
61. Smania, N., Aglioti, S. M., Girardi, F., Tinazzi, M., Fiaschi, A., Cosentino, A., & Corato, E. (2006). Rehabilitation of limb apraxia improves daily life activities in patients with stroke. *Neurology, 67*, 2050–2052.
62. van der Gaag, A., & Brooks, R. (2008). Economic aspects of a therapy and support service for people with long-term stroke and aphasia. *International Journal of Language & Communication Disorders / Royal College of Speech & Language Therapists, 43*(3), 233–244.
63. Vitali, P., Abutalebi, J., Tettamanti, M., Danna, M., Ansaldo, A. I., Perani, D., ... Cappa, S. F. (2007). Training-induced brain remapping in chronic aphasia: A pilot study. *Neurorehabilitation and Neural Repair, 21*(2), 152–160.
64. Wisenburn, B., & Mahoney, K. (2009). A meta-analysis of word-finding treatments for aphasia. *Aphasiology, 23*, 1338–1352.

Reason for exclusion: no original data

65. Crinion, J. T., & Leff, A. P. (2007). Recovery and treatment of aphasia after stroke: Functional imaging studies. *Current Opinion in Neurology, 20*, 667–673.
66. Henry, M. L., Beeson, P. M., & Rapcsak, S. Z. (2008). Treatment for lexical retrieval in progressive aphasia. *Aphasiology, 22*, 826–838.
67. Kelly, H., Brady, M. C., & Enderby, P. (2010). Speech and language therapy for aphasia following stroke. *Cochrane Database of Systematic Reviews, 5*, CD000425.
68. Kent, T. A., Rutherford, D. G., Breier, J. I., & Papanicolaou, A. C. (2009). What is the evidence for use dependent learning after stroke? *Stroke, 40*(3, Suppl. 1), S139–S140.

69. Meinzer, M., & Breitenstein, C. (2008). Functional imaging studies of treatment-induced recovery in chronic aphasia. *Aphasiology*, *22*, 1251–1268.
70. Meinzer, M., Elbert, T., Djundja, D., Taub, E., & Rockstroh, B. (2007). Extending the constraint-induced movement therapy (CIMT) approach to cognitive functions: Constraint-induced aphasia therapy (CIAT) of chronic aphasia. *NeuroRehabilitation*, *22*, 311–318.
71. Naeser, M. A., Martin, P. I., Treglia, E., Ho, M., Baker, E., & Kaplan, E. (2009). *Improved action naming in a severe, nonfluent aphasia patient following transcranial magnetic stimulation plus constraint-induced language therapy*. Poster presented at the 47th annual meeting of the Academy of Aphasia, Turku, Finland. Available from http://academy.angularis.org/Annual_Meeting/2009/CoverSheet/Site/Program/poster2.pdf.
72. Poslawsky, I. E., Schuurmans, M. J., Lindeman, E., & Hafsteinsdóttir, T. B. (2010). A systematic review of nursing rehabilitation of stroke patients with aphasia. *Journal of Clinical Nursing*, *19*(1–2), 17–32.
73. Pulvermüller, F., & Berthier, M. L. (2008). Aphasia therapy on a neuroscience basis. *Aphasiology*, *22*, 563–599.
74. Teasell, R., Foley, N. C., Salter, K., Bhogal, S. K., Jutai, J., & Speechley, M. (2009). *Evidence-based review of stroke rehabilitation*. Toronto, Ontario, Canada: Heart and Stroke Foundation of Ontario.

Reason for exclusion: included in original Cherney et al. (2008) review

75. Maher, L. M., Kendall, D., Swearingin, J. A., Rodriguez, A., Leon, S. A., Pingel, K., ... Gonzalez Rothi, L. J. (2006). A pilot study of use-dependent learning in the context of constraint induced language therapy. *Journal of the International Neuropsychological Society*, *12*, 843–852.

Reason for exclusion: non-English or non-peer-reviewed journal

76. Bilda, K., Matzner, K., Jochims, H., Breitenstein, C., Bildat, L., & Müller-Dohm, B. (2008). [Video-based conversation training: A study on high frequency learning of everyday dialogues in aphasia]. *Forum Logopadie*, *22*(5), 22–27.
77. Breitenstein, C., Kramer, K., Meinzer, M., Baumgartner, A., Floel, A., & Knecht, S. (2009). [Intense language training for aphasia. Contribution of cognitive factors]. *Nervenarzt*, *80*(2), 149–50.
78. Cherney, L. B. E., Cole, R., Van Vuuren, S., Hurwitz, R., & Ngampatipatpong, N. (2006). Computer treatment for aphasia: Efficacy and treatment intensity. *Archives of Physical Medicine and Rehabilitation*, *87*(10), E5.
79. Godecke, E., West, D., Cartwright, J., Cream, A., Ciccone, N., Hankey, G., ... Phillips, M. (2009). Intensive aphasia therapy in the early poststroke recovery phase: Is group intervention a viable therapy option? *International Journal of Stroke*, *4*(Suppl. 1), 28.

80. Jackson, C., Brown, J., Richardson, E., & Yaras, S. (2008, November). *Effects of constraint-induced aphasia therapy on discourse structure*. Paper presented at the annual convention of the American Speech-Language-Hearing Association, Chicago.
81. Krüger, S., Breunig, V., & Werner, R. (2009). [CIAT in clinical everyday life—results and practical experience]. *Forum Logopädie*, 23(6), 20–25.
82. Lauterbach, M., Leal, G., Aguiar, M., Fonseca, I., Farrajota, L., Fonseca, J., et al. (2007, September 10–12). *Intensive vs conventional speech therapy in aphasia due to ischaemic stroke: A randomized controlled trial*. Paper presented at the British Aphasiology Society 2007 Biennial Conference, Edinburgh, Scotland.
83. Lee, J. B., & Cherney, L. R. (2008). The changing "face" of aphasia therapy. *Perspectives on Neurophysiology and Neurogenic Speech and Language Disorders*, 18(1), 15–23.
84. Piron, L., Meneghello, F., De Pellegrin, S., Cremasco, A., & Bevilacqua, F. (2006). Early and intensive speech therapy: Follow-up and group control study on severe aphasia. *Neurorehabilitation and Neural Repair*, 20(1), 178.
85. Raymer, A. M., Maher, L. M., Patterson, J., & Cherney, L. (2007). Neuroplasticity and aphasia: Lessons from constraint-induced language therapy. *Perspectives on Neurophysiology and Neurogenic Speech and Language Disorders*, 17(2), 12–17.
86. Sun, H. F., Ni, C. M., Chaimay, B., Thinkhamrop, B., Thinkhamrop, J., Millar, D., et al. (2006). [Rehabilitative treatment of aphasia: Study and application]. *Chinese Journal of Clinical Rehabilitation*, 10, 125–127.
87. Vauth, F., Hampel, P., Scibor, M., Handschu, R., Richter, J., & Keidel, M. (2008). [Synchronic telepractice: A new (additional) form of aphasia therapy]. *Forum Logopädie*, 22(4), 12–19.
88. Virion, C. R. (2008). *"Go aphasia!": Examining the efficacy of constraint-induced language therapy for individuals with agrammatic aphasia*. University of Maryland, College Park, College Park, MD.
89. Wilssens, I. M. E. (2009). *Constraint induced aphasia therapy or semantic treatment in chronic fluent aphasia: A review and an explorative study*. Universiteit Utrecht, Utrecht, Netherlands.

Reason for exclusion: not population under review

90. Barnes, D. E., Yaffe, K., Belfor, N., Jagust, W. J., DeCarli, C., Reed, B. R., & Kramer, J. H. (2009). Computer-based cognitive training for mild cognitive impairment: Results from a pilot randomized, controlled trial. *Alzheimer Disease and Associated Disorders*, 23(3), 205–210.
91. Breitenstein, C., Zwitserlood, P., de Vries, M. H., Feldhues, C., Knecht, S., & Dobel, C. (2007). Five days versus a lifetime: Intense associative vocabulary training generates lexically integrated words. *Restorative Neurology and Neuroscience*, 25, 493–500.
92. Cherney, L. R., & Halper, A. S. (2008). Novel technology for treating individuals with aphasia and concomitant cognitive deficits. *Topics in Stroke Rehabilitation*, 15, 542–554.

93. Henry, M. L., Beeson, P. M., & Rapcsak, S. Z. (2008). Treatment for anomia in semantic dementia. *Seminars in Speech and Language, 29*(1), 60–70.
94. Rogalski, Y., & Edmonds, L. A. (2008). Attentive reading and constrained summarisation (ARCS) treatment in primary progressive aphasia: A case study. *Aphasiology, 22*, 763–775.

Reason for exclusion: not a study or systematic review with original data

95. Balardin, J. B., & Miotto, E. C. (2009). A review of constraint-induced therapy applied to aphasia rehabilitation in stroke patients. *Dementia & Neuropsychologia, 3*(4), 275–282.
96. Ball, A. L., Grether, S., Szaflarski, J. P., Griffith, N., Al-Fwawes, F., Newmeyer, A. J., ... Reichhardt, R. (2006). Constraint-induced aphasia treatment (CIAT) in patients with chronic language impairment after stroke. *Neurology, 66*, A324–A325.
97. Berthier, M. L., Green, C., Lara, J. P., Higuera, C., Barbancho, M. A., Davila, G., et al. (2008). Awakening the language cortex with memantine and constraint-induced speech-language therapy: A randomised placebo-controlled trial in chronic post-stroke aphasia. *European Journal of Neurology, 15*, 28.
98. Berthier, M. L., Green, C., Lara, P., Higuera, C., Barbancho, M. A., Davila, G., Pulvermüller, F. (2009). Memantine and constraint-induced aphasia therapy in chronic post-stroke aphasia: A randomized controlled trial with an open-label extension phase. *Neurology, 72*, A481–A482.
99. Dembowski, J. S. (2009). Available studies fail to disambiguate contributions of language modality constraint and treatment intensity to effects of constraint-induced language therapy. *Evidence-Based Communication Assessment & Intervention, 3*(2), 72–78.
100. Kiran, S., & Bassetto, G. (2008). Evaluating the effectiveness of semantic-based treatment for naming deficits in aphasia: what works? *Seminars in Speech and Language, 29*(1), 71–82.
101. Lara, J. P., Barbancho, M. A., Berthier, M. L., Green, C., Navas, P., Dawid-Milner, M. S., ... Gonzalez-Baron, S. (2009). ERPs correlates of recovery from chronic post-stroke aphasia in patients treated with memantine and constraint-induced aphasia therapy. *European Journal of Neurology, 16*, 457–457.
102. Lillie, R., & Mateer, C. A. (2006). Constraint-based therapies as a proposed model for cognitive rehabilitation. *Journal of Head Trauma Rehabilitation, 21*(2), 119–130.
103. Marshall, R. C. (2008). The impact of intensity of aphasia therapy on recovery. *Stroke, 39*(2), e48; author reply e49.
104. Raymer, A. M., Beeson, P., Holland, A., Kendall, D., Maher, L. M., Martin, N., ... Gonzalez Rothi, L. J. (2008). Translational research in aphasia: From neuroscience to neurorehabilitation. *Journal of Speech, Language, and Hearing Research, 51*, S259–S275.
105. Taub, E., Uswatte, G., Mark, V. W., & Morris, D. M. (2006). The learned nonuse phenomenon: Implications for rehabilitation. *Europa Medicophysica, 42*(3), 241–256.
106. Thompson, C. K., & den Ouden, D.-B. (2008). Neuroimaging and recovery of language in aphasia. *Current Neurology and Neuroscience Reports, 8*, 475–483.

107. Wilson, B. A. (2010). Brain injury: recovery and rehabilitation. *Wiley Interdisciplinary Reviews: Cognitive Science*, 1(1), 108–118.